

# Radiation pressure forces and blowout sizes for particles in debris disks

Jessica A. Arnold<sup>a,\*</sup>, Alycia J. Weinberger<sup>a</sup>, and Gorden Videen<sup>b,c</sup>

<sup>a</sup> *Department of Terrestrial Magnetism, Carnegie Institution for Science, 5421 Broad Branch Rd., Washington, DC 20015, USA*

<sup>b</sup> *US Army Research Laboratory RDRL-CIE-S, 2800 Powder Mill Rd., Adelphi, MD 20783, USA*

<sup>c</sup> *Space Science Institute, 4750 Walnut Street, Boulder Suite 205, CO 80301, USA*

*\*Presenting author (jarnold@carnegiescience.edu)*

Debris disks are a type of circumstellar disk that contain dust generated by collisions and disruptions of protoplanets and/or planetesimals. To interpret scattered light observations of debris disks, it is useful to compare modeled grain size distributions to radiation pressure blowout sizes. The ratio of radiation pressure to gravitational forces ( $\beta$ ) acting on a dust grain depends on grain composition, size, and structure. Typically,  $\beta$  is calculated using the assumption of compact, spherical particles or accounting for porosity via the Maxwell-Garnett mixing rule (e.g., [1]). Calculations of radiation pressure balance for porous, irregular dust grains have been carried out for a handful of cases [2–4] using the discrete dipole approximation (DDA) method [5]. However, due to computational considerations, these focused on submicron particles that only require a small number of dipoles ( $N \leq 2048$ ), but may be below the blowout size of some systems.

Here we present comparisons between Mie, Maxwell-Garnett, and DDA calculations of  $\beta$  for micron-sized grains using different stellar luminosities and grain compositions. The grain shapes and DDA implementation used to generate scattering and absorption efficiencies are similar to [6]. Stellar properties were chosen to correspond to stars known to host debris disks.

## References

- [1] Mukai, T., H. Ishimoto, T. Kozasa, *et al.*, 1992: Radiation pressure forces of fluffy porous grains. *Astron. Astrophys.* **262**, 315–320.
- [2] Kimura, H., H. Okamoto, and T. Mukai, 2002: Radiation pressure and the Poynting–Robertson effect for fluffy dust particles. *Icarus* **157**, 349–361.
- [3] Kirchsclanger, F., and S. Wolf, 2013: Porous dust grains in debris disks. *Astron. Astrophys.* **552**, A54.
- [4] Silsbee, K., and B. Draine, 2016: Radiation pressure on fluffy submicron-sized grains. *Astrophys. J.* **818**, 2.
- [5] Draine, B. T., and P. J. Flatau, 1994: Discrete dipole approximation for scattering calculations. *J. Opt. Soc. Am. A* **11**, 1491–1499.
- [6] Zubko, E., D. Petrov, Y. Shkuratov, *et al.*, 2005: Discrete dipole approximation simulations of scattering by particles with hierarchical structure. *Appl. Opt.*, **44**, 6479–6485.

Preferred mode of presentation: Oral/Poster